





Injuries can damage soft tissues such as muscles, ligaments and tendons. When soft tissue is injured blood vessels are usually damaged too. This leads to bleeding and swelling around the damaged tissue which can cause pain, further damage, impaired healing and loss of joint range of motion. Therefore reducing bleeding at the site of injury is vital. The first 48 hours following injury are the most important in the management of acute soft tissue injuries. Appropriate treatment will help to relieve immediate symptoms and speed up your recovery. The initial treatment should follow the RICED (Rest, Ice, Compression, Elevation, Diagnosis) regime and avoid HARM-ful (Heat, Alcohol, Running, Massage) activities.

<p>REST</p> <p>Rest reduces further injury, bleeding and swelling. Stop activities as soon as your injury occurs and avoid movement of the injured area when possible.</p>	 <p>COMPRESSION</p>	<p>COMPRESSION</p> <p>Compression of the injured area reduces bleeding and swelling. This can be achieved by wrapping the area in a crepe bandage that extends above and below the injured area.</p>
<p>ICE</p> <p>Ice reduces pain, spasm, bleeding and swelling. Apply for 20 minutes every 2 hours for the first 48 hours. This can be achieved with ice cubes wrapped in a moist towel, placed around the injured area and held in place with a bandage.</p>	 <p>REST</p> <p>ELEVATION</p>	<p>ELEVATION</p> <p>Elevation of the injury above the level of the heart reduces pain, bleeding and swelling. For lower limb injuries, this can be achieved by lying down with your affected leg resting on a pillow or chair. For upper limb injuries, you can wear a sling.</p>
<p>Alternatively, water frozen in a foam cup will produce a large ice block that can be moved across the skin of the injured area (apply a little oil to the skin first). Other methods are immersion in icy water and cold gel packs.</p>	 <p>ICE MASSAGE</p> <p>ICE PACK</p>	<p>DIAGNOSIS</p> <p>Consult a medical professional as soon as you are able for expert advice on the severity of the injury and the necessary treatment.</p>
<p>DIAGNOSIS</p>	 <p>DIAGNOSIS</p>	<p>DIAGNOSIS</p> <p>Consult a medical professional as soon as you are able for expert advice on the severity of the injury and the necessary treatment.</p>

Avoid HARM-ful activities:

HEAT: Heat increases bleeding at the injured area. Avoid hot baths, hot showers, heat packs, heat rubs/liniments and hot water bottles for at least 72 hours after injury.

ALCOHOL: Alcohol increases bleeding and swelling at the injured area and can lengthen recovery time.

RUNNING: Running or exercise of the injured area can cause further damage within the first 72 hours after injury.

MASSAGE: Massage increases bleeding and swelling at the injured area and should also be avoided in the first 72 hours post-injury.